

Child's Name: _____ Birthdate: _____ Male/Female School: _____
Last, First month/day/year

Address _____ Phone: _____ Grade: _____
Street City Zip

Santa Clara County Public Health Department TB Risk Assessment for School Entry

This form must be completed by a licensed health professional and returned to the child's school.

1. Was your child born in Africa, Asia, Latin America, or Eastern Europe? Yes No
2. Has your child traveled to a country with a high TB rate* (for more than a week)? Yes No
3. Has your child been exposed to anyone with tuberculosis (TB) disease? Yes No
4. Has a family member or someone your child has been in contact with had a positive TB test or received medications for TB? Yes No
5. Was a parent, household member or someone your child has been in close contact with, born in or traveled to a country with a high TB rate? Yes No
6. Has another risk factor for TB (i.e. one of those listed on the back of this page)? Yes No

* This includes countries in Africa, Asia, Latin America or Eastern Europe. For travel, the risk of TB exposure is higher if a child stayed with friends or family members for a cumulative total of 1 week or more.

If YES, to any of the above, the child has an increased risk of TB infection and should have a TST/ IGRA.

All children with a positive TST/IGRA result must have a medical evaluation, including a chest X-ray. Treatment for latent TB infection should be initiated if the chest X-ray is normal and there are no signs of active TB. If testing was done, please attach or enter results below.

Tuberculin Skin Test (TST/Mantoux/PPD) Date given: _____ Date read: _____	Induration _____ mm Impression: <input type="checkbox"/> Negative <input type="checkbox"/> Positive
Interferon Gamma Release Assay (IGRA) Date: _____	Impression: <input type="checkbox"/> Negative <input type="checkbox"/> Positive <input type="checkbox"/> Indeterminate
Chest X-Ray (required with positive TST or IGRA) Date: _____	Impression: <input type="checkbox"/> Normal <input type="checkbox"/> Abnormal finding
<input type="checkbox"/> LTBI treatment (Rx & start date): _____	<input type="checkbox"/> Prior TB/LTBI treatment (Rx & duration): _____
<input type="checkbox"/> Contraindications to INH or rifampin for LTBI	<input type="checkbox"/> Offered but refused LTBI treatment

Providers, please check one of the boxes below and sign:	
<input type="checkbox"/> Child has no TB symptoms, none of the above or other risk factors for TB and does not require a TB test.	
<input type="checkbox"/> Child has a risk factor, has been evaluated for TB and is free of active TB disease.	
_____	_____
Health Provider Signature, Title	Date

Name/Title of Health Provider:

Facility/Address:

Phone number:

Fax number:

County of Santa Clara

Public Health Department



Tuberculosis Prevention & Control Program
976 Lenzen Avenue, Suite 1700
San José, CA 95126
408.885.2440

Risk Factors for Tuberculosis (TB) in Children

- Have clinical evidence or symptoms of TB
- Have a family member or contacts with history of confirmed or suspected TB
- Are in foreign-born families from TB endemic countries (including countries in Africa, Asia, Latin America or Eastern Europe)
- Travel to countries with high rate of TB
- Contact with individual(s) with a positive TB test
- Abnormalities on chest X-ray suggestive of TB
- Adopted from any high-risk area or live in out-of-home placements
- Live with an adult who has been incarcerated in the last five years
- Live among or frequently exposed to individuals who are homeless, migrant farm workers, residents of nursing homes, or users of street drugs
- Drink raw milk or eat unpasteurized cheese (i.e. queso fresco or unpasteurized cheese)
- Have, or are suspected to have, HIV infection or live with an adult with HIV seropositivity. See below for testing methods in children with HIV or other immunocompromised conditions.

Testing Methods

A Mantoux tuberculin skin test (TST) or an Interferon Gamma Release Assay (IGRA) (for children aged 4 and older) should be used to test those at increased risk. A TST of ≥ 10 mm is considered positive. If a child has had contact with someone with active TB (yes to question 3 on reverse) then TST ≥ 5 mm is considered positive.

Screening should be performed by CXR in addition to a TST/IGRA (consider doing both) and symptom review in HIV infected or suspected HIV, other immunocompromised conditions or if a child is taking immunosuppressive medications such as prednisone or TNF-alpha antagonists.

Referral, Treatment, and Follow-up of Children with Positive TB Tests

- All children with a positive TST or IGRA result should have a medical evaluation, including a chest X-ray.
- Report any confirmed or suspected case of TB disease to the TB Control Program within 1 day, including any child with an abnormal chest X-ray.
- If TB disease is not found, treat children and adolescents with a positive TST or IGRA for latent TB infection (LTBI).
- Isoniazid (INH) is the drug of choice for the treatment of LTBI in children and adolescents. The length of treatment is 9 months with daily dosing: 10-15mg/kg (maximum 300 mg).
- For management and treatment guidelines for TB or LTBI, go to: www.cdc.gov/tb or contact the TB Control Program at (408) 885-4214.

References

American Academy of Pediatrics, Committee on Infectious Diseases. Tuberculosis. In L.K. Pickering (Ed.), 2009 *Red Book: Report of the Committee on Infectious Diseases*. 27th ed. El Grove Village, IL: American Academy of Pediatrics, 2009:680-701.

California Health and Safety Code Section 121515.

Pediatric Tuberculosis Collaborative Group. Targeted Tuberculin Skin Testing and Treatment of Latent Tuberculosis Infection in Children and Adolescents. *Pediatrics* 2004; 114 (14):1175-1201.

Pang J, Teeter LD, Katz DJ, et al. Epidemiology of Tuberculosis in Young Children in the United States. *Pediatrics*, 2014:494-504.

Board of Supervisors: Mike Wasserman, Cindy Chavez, Dave Cortese, Ken Yeager, S. Joseph Simitian,
County Executive: Jeffrey V. Smith